



 <h1>Choosing Wisely®</h1> <p><i>An initiative of the ABIM Foundation</i></p>	  <p>AMERICAN ACADEMY OF FAMILY PHYSICIANS</p> 
--	---

Bone-density tests

When you need them—and when you don't


It's worth getting a bone-density test if you're older or have other risk factors for weak bones, because without it the first sign of osteoporosis is usually a broken bone. But if you're not at higher risk, you should think twice about the test. Here's why.

The test usually isn't necessary in younger people without risk factors for weak bones.

Many women and some men are routinely screened for weak bones with an imaging test called a dual-energy X-ray absorptiometry (DEXA) scan. If it shows that you have outright osteoporosis, the results can help you and your doctor decide how to treat the problem, usually with drugs. But many people learn they have only mild bone loss, a condition known as osteopenia, and for them the risk of fracture is often quite low.

It can pose risks.

A DEXA scan isn't likely to harm you directly. But a diagnosis of osteopenia can lead to treatment with such drugs as alendronate (Fosamax and generic), ibandronate (Boniva and generic), and risendronate (Actonel, Atelvia, and gener-



ic), which pose numerous risks. Those include thigh fractures; throat or chest pain; difficulty swallowing; heartburn; and more rarely, bone, eye, joint, and muscle pain; bone loss in the jaw; and possibly abnormal heart rhythm. In addition, there is little evidence that people with osteopenia get much benefit from the drugs. Other types of osteoporosis drugs are linked to risks

such as blood clots, heart attacks, strokes, and serious infections.

It can be a waste of money.

A DEXA scan costs about \$132, according to HealthcareBlueBook.com. Though that isn't as expensive as some tests, any money spent on unnecessary tests is money wasted. In addition, a month's supply of generic alendronate costs \$38 to \$70. Fosamax, the brand-name version, costs \$125 to \$148. People often take the drugs for years and sometimes indefinitely.

So when is the test warranted?

Women should have their bone density measured at age 65. Men 70 and older might also want to talk with their doctor about the test. Women younger than 65 and men 50 to 69 should consider it if they have risk factors such as a fracture from minor trauma, rheumatoid arthritis, a parent who had a hip fracture, or a history of smoking, heavy drinking, or long-term use of corticosteroid drugs. Whether you need a follow-up bone-density test depends on the results of the initial scan.

USING THIS INFORMATION

This information is provided for you to use in discussions with your health-care provider. The content is for educational use only and is not a substitute for professional medical advice, diagnosis, or treatment. Unfortunately, we cannot answer individual medical questions. Always seek the advice of a physician or other qualified health-care provider with questions about any medical condition. Never disregard, avoid, or delay in obtaining medical advice from a health-care provider because of something you have read in this report. Use of this report is at your own risk. Consumer Reports, AAFP, the ABIM Foundation, Archives of Internal Medicine, and their distributors are not liable for any loss, injury, or other damage related to your use of this report.

The report is intended solely for consumers' personal, noncommercial use and may not be altered or modified in any way or used in advertising, for promotion, or for any other commercial purpose. Special permission is granted to organizations in the Consumer Reports health communication program to disseminate free copies of this report in print or digital (PDF) formats to individual members and employees. Learn more at ConsumerHealthChoices.org or send an e-mail to HealthImpact@cr.consumer.org.

Published by Consumer Reports © 2012 Consumers Union of U.S., Inc., 101 Truman Ave., Yonkers, NY 10703-1057. Developed in cooperation with AAFP. Portions are derived from AAFP's "Five Things Physicians and Patients Should Question" list. © 2012 AAFP. This list is based on the "The Top 5 Lists in Primary Care: Meeting the Responsibility of Professionalism," the Good Stewardship Working Group, *Arch Intern Med.* 2011;171:1385-1390. "Top 5 Lists in Primary Care" were first published as part of the Less is More™ series in the Archives of Internal Medicine. © 2011 American Medical Association.

CONSUMER REPORTS' Advice

How can you keep your bones strong?

The following steps can help protect against fractures by building bone:

- **Do weight-bearing exercises.** Aim for at least 30 minutes a day of activities such as walking and weightlifting.

- **Get enough calcium and vitamin D.**

Aim for at least 1,200 milligrams of calcium a day, from green leafy vegetables, low-fat dairy products, canned sardines or salmon, and if necessary, supplements. Consider a vitamin D supplement if



you are a woman after menopause or get little sun exposure. Aim for 600 international units a day, or 800 IU if you are 70 or older.

- **Avoid smoking and excessive alcohol use.** Try a smoking-cessation program or nicotine-replacement product, and limit yourself to one drink a day for women, two for men.

- **Minimize bone-sapping medication.**

Talk with your doctor about alternatives to drugs such as corticosteroids, proton pump inhibitors used to treat heartburn, and certain newer antidepressants.

- **Reduce your risk of falling.** Eliminate small rugs and loose extension cords in your home, use brighter lights, and install grab bars and use rubber mats in the bathroom. Have your vision checked, wear nonslip shoes, and take classes that increase balance and strength, such as tai chi and yoga. Ask your doctor whether any drug you take could impair your balance.